

Mercury's formation impact splattered Earth with material (Forwarded)

Source: <http://sci.tech-archive.net/Archive/sci.astro/2006-04/msg00057.html>

- *From:* Andrew Yee <"ayee\"@nova.astro.utoronto.ca">
 - *Date:* Wed, 05 Apr 2006 21:35:24 -0400
-

ROYAL ASTRONOMICAL SOCIETY PRESS INFORMATION NOTE

Issued by RAS Communications Officers:

Peter Bond

Tel: +44 (0)1483-268672 Fax: +44 (0)1483-274047

AND

Anita Heward

Tel: +44 (0)1483-420904

NATIONAL ASTRONOMY MEETING PRESS ROOM (4 - 7 APRIL ONLY):

Tel: +44 (0)116-229-7474 or 229-7475 or 252-3312 or 252-3531

Fax: +44 (0)116-252-3531

RAS Web site:

<http://www.ras.org.uk/>

RAS National Astronomy Meeting web site:

<http://www.nam2006.le.ac.uk/index.shtml>

CONTACT DETAILS ARE LISTED AT THE END OF THIS RELEASE.

EMBARGOED FOR 02:01 BST, WEDNESDAY, 5 APRIL 2006 (01:01 GMT, WEDNESDAY 5 APRIL)

Ref.: PN 06/17 (NAM 10)

MERCURY'S FORMATION IMPACT SPLATTERED EARTH WITH MATERIAL

New computer simulations of Mercury's formation show the fate of material blasted out into space when a large proto-planet collided with a giant asteroid 4.5 billion years ago. The simulations, which track the material over several million years, shed light on why Mercury is denser than expected and show that some of the ejected material would have found its way to the Earth and Venus.

"Mercury is an unusually dense planet, which suggests that it

Mercury's formation impact splattered Earth with material (Forwarded)

contains far more metal than would be expected for a planet of its size. We think that Mercury was created from a larger parent body that was involved in a catastrophic collision, but until these simulations we were not sure why so little of the planet's outer layers were reaccreted following the impact," said Dr Jonti Horner, who is presenting results at the Royal Astronomical Society's National Astronomy Meeting on 5th April.

To solve this problem, Dr Horner and his colleagues from the University of Bern ran two sets of large-scale computer simulations. The first examined the behaviour of the material in both the proto-planet and the incoming projectile; these simulations were among the most detailed to date, following a huge number of particles and realistically modelling the behaviour of different materials inside the two bodies. At the end of the first simulations, a dense Mercury-like body remained along with a large swathe of rapidly escaping debris. The trajectories of the ejected particles were then fed in to a second set of simulations that followed the motion of the debris for several million years. Ejected particles were tracked until either they landed on a planet, were thrown into interstellar space, or fell into the Sun. The results allowed the group to work out how much material would have fallen back onto Mercury and investigate other ways in which debris is cleared up in the Solar System.

The group found that the fate of the debris depended on whereabouts Mercury was hit, both in terms of its orbital position and in terms of the angle of the collision.

Whilst purely gravitational theory suggested that a large fraction of the debris would eventually fall back onto Mercury, the simulations showed that it would take up to 4 million years for 50% of the particles to land back on the planet and in this time many would be carried away by solar radiation. This explains why Mercury retained a much smaller proportion than expected of the material in its outer layers.

The simulations also showed that some of the ejected material made its way to Venus and the Earth. While this is only a small fraction, it illustrates that material can be transferred between the inner planets relatively easily. Given the amount of material that would have been ejected in such a catastrophe, it is likely that there is a reasonable amount (possibly as much as 16 million billion tonnes [1.65×10^{19} kg]) of proto-Mercury in the Earth.

IMAGES

http://www.ras.org.uk/images/stories/press/protomercury_formation1.jpg (46KB)
http://www.ras.org.uk/images/stories/press/protomercury_formation2.jpg (65KB)
http://www.ras.org.uk/images/stories/press/protomercury_formation3.jpg (55KB)

Mercury's formation impact splattered Earth with material (Forwarded)

Caption: The images show the evolution of the impact over the first three hours from the time of collision. The red colour on the plots shows the 'core' material (iron), while the blue shows the lighter mantle material (silicates, modelled as a rock type called dunite).

© Horner et Al 2006

NOTES FOR EDITORS

The 2006 RAS National Astronomy Meeting is hosted by the University of Leicester. It is sponsored by the Royal Astronomical Society, the UK Particle Physics and Astronomy Research Council (PPARC), the University of Leicester and the National Space Centre, Leicester.

CONTACTS

Dr. Jonathan Horner
Physikalisches Institut
Space Research & Planetary Sciences
Sidlerstrasse 5, 3012 Bern, Switzerland
Tel: ++41 (0) 31 631 8546
Fax: ++41 (0) 31 631 4405
Web: <http://www.phim.unibe.ch/~horner/>

From Tuesday 4th – Wednesday 5th April, Dr Horner can be contacted via the NAM press office (see above).