

Re: Sub clinical Cushing's syndrome

Source: <http://sci.tech-archive.net/Archive/sci.med.cardiology/2007-06/msg00717.html>

- *From:* tauzinger <tauzinger@xxxxxxxx>
 - *Date:* Fri, 29 Jun 2007 19:15:04 -0000
-

On Jun 29, 2:59 pm, Susan <neverm...@xxxxxxxx> wrote:

x-no-archive: yes

<http://www.csrf.net/Articles2.htm#Treatment>

"What populations are most likely to have subclinical Cushing's syndrome? One might look for SCS in two main patient populations with the expectation that making this diagnosis would lead to a beneficial therapeutic intervention: 1) patients with an incidentally discovered tumor that may inappropriately secrete cortisol (adrenal gland) or ACTH (pituitary gland). Usually these lesions are called incidentalomas. 2) patients that display non-specific pathologies that may be secondary to hypercortisolism such as obesity, diabetes, hypertension, osteoporosis etc.."

Susan

<http://endocrine.niddk.nih.gov/pubs/cushings/cushings.htm>
says "10-15 million a year", but it doesn't say if this is US or worldwide.

http://en.wikipedia.org/wiki/Cushing's_syndrome

"Incidence of causes of Cushing's syndrome per million person-years:

- * Cushing's disease: 10
- * Primary adrenal: 3
- * Ectopic ACTH: 1.4"

Thanks,

Thomas

<http://ecg.auzinger.org>

.